

# ***INFECTION CONTROL***

## ***TEST***

- 1. What are some ways to contact the IC staff?**
  - a. Phone
  - b. Email
  - c. Fax
  - d. All of the above
- 2. You have sustained a needle stick injuries. What should you do?**
  - a. Perform first aid by cleaning or flushing the exposed area, report the exposure to your supervisor ASAP, report to WAMC-ED for evaluation, treatment, and counseling; and follow up with the Occupational Health Clinic within 1 business day.
  - b. Notify your supervisor, and report to the occupational health located in the wellness clinic.
  - c. Wash area with soap/water, report to the emergency department, complete paperwork, and notify your supervisor.
  - d. Wash area with soap/water, complete paperwork and report to occupational health.
- 3. All personnel must:**
  - a. Be included in the exposure documentation, receive Hep B vaccine, and receive annual BBP training.
  - b. Be excluded in the exposure documentation, receive Hep A vaccine, and receive annual Flu training.
  - c. Receive bi-annual BBP training
  - d. Receive Hep B vaccine annually
- 4. Which is not correct when cleaning a blood spill:**
  - a. Contain spill, wear PPE, wipe up the spill with paper towels, (throw away in red bag) wash area with hospital approved disinfectant, remove PPE, wash hands
  - b. contain spill, notify housekeeping
  - c. contain spill, wipe the blood with a cloth towel, throw cloth towel in red bag, wash area with soap and water
- 5. The Occupational Exposure Control Plan is:**
  - a. A complete list of employees by name and job description at Womack. This file is maintained in Civilian personnel office/military personnel office.
  - b. Policy that details measures at Womack and employees will take to decrease risk of blood borne pathogens exposure. Policy located in the infection control manual.

- c. MEDCEN memo that outlines the process to follow in the event that an exposure occurs and WAMCs responsibility to prevent and exposure if one does occur.
- 6. Personal protective equipment PPE can include all of the following except:**
- a. Prescription eye glasses
  - b. gloves
  - c. Protective clothing like gowns and bunny suits
  - d. Masks, goggles
- 7. The following items are considered RMW (Regulated Medical Waste) and are disposed of in a red biohazard bag.**
- a. Gowns, gloves, masks used while caring for a patient on contact or droplet or airborne precautions for (MRSA, chickenpox, bacterial meningitis, tuberculosis)
  - b. Animal waste, specimen transport bags, unused red bags, bloody gauze, chest tubes
  - c. diapers, sanitary pads, urine, feces
- 8. Symptoms of pulmonary tuberculosis (TB) include:**
- a. weight loss
  - b. productive cough more than three weeks
  - c. night sweats, fever, chills
  - d. all of the above
- 9. In the event a patient present to your clinic with symptoms of tuberculosis (TB) you should?**
- a. Place a surgical mask on you on the patient
  - b. isolate patient (removed from waiting room)
  - c. notify infection control
  - d. all of the above
- 10. Which type of precautions requires all employees to treat all body fluids, secretions, and tissues of All patients as potentially infectious? (wet and not yours)**
- a. Contact
  - b. Droplet
  - c. Standard
  - d. Airborne
- 11. What is the most important way to reduce the transmission of infectious agents by staff to patients?**
- a. Stay away from sick people
  - b. wearing gloves
  - c. perform hand hygiene
  - d. none of the above
- 12. The WHO's 5 Moments for Hand Hygiene at point of care:**
- a. Before patient contact, before an aseptic task, after body fluid exposure risk, after patient contact, and after contact with the patient's surroundings
  - b. Before patient contact, before an aseptic task, after body fluid exposure risk, after patient contact, and after using the bathroom
  - c. Before patient contact, before an aseptic task, after body fluid exposure risk, after patient contact, and after eating

- d. Before an aseptic task, after body fluid exposure risk, after patient contact, and after contact with the patient's surroundings

**13. Employees who wear prescription eye glasses do not have to wear goggles/protection?**

- a. True
- b. False

**14. Examples of illnesses that require airborne precaution are?**

- a. Tuberculosis (TB), measles, varicella, disseminated shingles
- b. meningitis
- c. influenza
- d. none of the above

**15. What is the contact time for Cavi-Wipes?**

- a. 1 minute
- b. 3 minutes
- c. 5 minutes
- d. 10 minutes

**16. What HAIs are preventable when utilizing aseptic technique?**

- a. CAUTI, CLABSI, SSI, C. diff
- b. VAP
- c. Bathing
- d. Emptying a urinary catheter

**17. Always pre-spike a bag of IV fluids 3 hours prior to use.**

- a. True
- b. False

**18. It is acceptable to use medications packaged as single-dose or single-use for more than one patient. This includes ampoules, bags, and bottles of intravenous solutions.**

- a. True
- b. False

**19. For an infection to be transmitted, all links of the Chain of Infection must occur.**

- a. True
- b. False

**20. When do sharps containers have to be emptied?**

- a. ¼ full
- b. ½ full
- c. ¾ full
- d. 2/3 full